



Babak Mohajer, MD
20 West 20th Street Suite 703
New York, NY 10011
T (212)260-6505
F (212)260-3060

Date of Procedure: _____ **Time: **** _____ ******

The procedure is done at **East Side Endoscopy Center (380 2nd Ave. & 22nd St.)**
(For more information on the facility please go to www.esecgi.com)
****please arrive 45 minutes**** earlier than your procedure time.

Please bring your insurance card and Photo ID with you

Starting ONE WEEK before the procedure

- Please make sure if you are on Plavix, Coumadin and any other blood thinners, Please discuss with Dr Mohajer a week prior to scheduling your procedure.
- If you cannot hold off on Plavix or Coumadin or any other blood thinners please inform Dr.Mohajer.
- If you are currently taking Blood Pressure medications, Please be sure to take them very early the morning of the procedure with a sip of water!!!
- If you take insulin please make sure you informed Dr. Mohajer at the time of your consultation/Follow-up visit. (not the day of your procedure)
- If you cancel and reschedule your colonoscopy and begin new medication please inform Dr.Mohajer.

IT'S A MUST THAT YOU HAVE SOMEONE PICK YOU UP.

You should have your escort pick you up approximately 1 ½-2 hour after your procedure starts. If you don't have any one please be aware they might cancel your procedure.

Any cancellation or rescheduling must be done with our office not the facility.

(If you must cancel please give 48hr notice prior. if no 48hr notice is given we will charge you a \$100.00 fee)

Any questions or concerns please feel free to call us at the above number.

!! Please note if you are taking recreational drugs (i.e. Cocaine) please inform the doctor, you must be off it 1 week prior to your procedure. If not they will cancel your procedure!!



Dr. Babak Mohajer
20 West 20th Street Suite 703
New York City, NY 10011
212-260-6505

CLEAR LIQUID DIET

The day before your procedure (Starting from the morning you will be on a clear liquid diet all day)

THIS DIET PROVIDES FLUIDS THAT LEAVE LITTLE RESIDUE AND ARE EASILY ABSORBED WITH MINIMAL DIGESTIVE ACTIVITY. THIS DIET IS INADEQUATE IN ALL ESSENTIAL NUTRIENTS AND IS RECOMMENDED ONLY IF CLEAR LIQUIDS ARE TEMPORARILY NEEDED. **NO DARKS COLORS SUCH AS RED OR PURPLE LIQUIDS SHOULD BE CONSUMED!**

DRINKS ALLOWED



- CARBONATED BEVERAGES (clear)
- FRUIT FLAVORED DRINKS (clear)
- APPLE JUICES
- WHITE GRAPE JUICE
- LEMONADE
- CLEAR BROTH (CONSOMME)
- CLEAR FLAVORED GELATIN, JELL-O (JELLO –LEMON OR LIME ,NO FRUIT OR TOPPINGS)
- POPSICLES (NO RED OR PURPLE)
- SUGAR, HONEY, SYRUP, CLEAR HARD CANDY, SALT
- GREEN TEA (please make sure it's not too dark)
- COCONUT WATER

All Day



FOODS NOT ALLOWED- NO SOLIDS AT ALL THE DAY BEFORE-Just Liquids

MILK, MILK DRINKS, FRUIT JUICES WITH UNSTRAINED FRUITS, ALCOHOL BEVERAGES, VEGETABLES, MEATS, GRAINS, STARCHES, NO COFFEE

*****Avoid Seeds / Nuts for at least 5 days prior to your procedure*****

Last CLEAR Liquid meal should be at 12midnight unless otherwise instructed.

BABAK MOHAJER, MD.
20 West 20th Street Suite 703
New York City, NY 10011
212-260-6505 P
212-260-3060 F

Tri-Lyte Instructions

Note: Stop aspirin or aspirin containing products one week prior to the exam. Please advise the office if you are taking any blood thinners (Coumadin, Plavix).

****Before beginning the following steps have your prescription for Tri-Lyte with flavor packs filled. **** (you should have it filled a few days before your Procedure)

The day before your procedure:

Be on a clear liquid diet. No solid foods the entire day before your procedure.

- 1.** In the morning, prepare your Tri-Lyte solution and refrigerate.
- 2.** Have a clear liquid breakfast
- 3.** Have a clear liquid lunch
- 4.** At 4pm you may have clear liquid dinner.
- 5.** At 6pm begin drinking the Tri-Lyte solution as instructed. Drink 8 oz. every 10-15 minutes.

*you will have consumed several glassfuls before having the first watery bowel movement.

*Initially, you may feel slightly bloated, but will become more comfortable as you continue to have bowel movements.

*It will take approximately 3 hours to finish the solution. Continue drinking the solution until your stool runs clear. Diarrhea will continue for about 1 hour after you finish drinking the solution.

****Do not eat anything until after your examination. ****

Any Questions please call us at the above number.